

Computational Statistics

Monte Carlo Markov Chains

1. Construct a random walk MH sampler to generate a sample of 10,000 observations from the Laplace distribution,

$$f(x) = \frac{1}{2}e^{-|x|}, \quad -\infty < x < +\infty.$$

Use $\epsilon \sim \mathcal{N}(0, \sigma^2)$ to generate proposals $y = x^{(t-1)} + \epsilon$. Compare the generated series for various values of σ^2 . Draw a histogram of the sampled values, together with the target density.

2. Consider the model

$$y_i = \beta x_i + u_i, \quad u_i \sim \mathcal{N}(0, 1), \quad i = 1, \dots, n,$$

with the gamma prior distribution $\beta \sim G(2, 1)$, $\beta > 0$.

- (a) Verify that the posterior distribution is

$$f(\beta|y_1, \dots, y_n) \propto \beta \exp(-\beta) \exp\left[-\frac{1}{2} \sum_{i=1}^n (y_i - \beta x_i)^2\right] \mathbb{1}_{(0, +\infty)}(\beta).$$

Note that this distribution does not have a standard form.

- (b) Construct an MH algorithm to sample from this distribution with an independence kernel, where the kernel is the prior distribution.
 - (c) Generate a dataset by choosing $n = 50$, x_i from $\mathcal{N}(0, 1)$ and a value of β from its prior distribution. Apply the MH algorithm to this dataset.
 - (d) Plot the sample path, the histogram of simulated values and the autocorrelation function (use function `acf`).
 - (e) Compute the posterior expectation of β and its standard error.
3. The dataset `coal.dat` is a time series of the numbers of disasters at coal mines annually between 1851 and 1962. We assume that the number of coal mining disasters follows a Poisson distribution $\mathcal{P}(\theta_1)$ until some unknown change point k , and then a Poisson distribution $\mathcal{P}(\theta_2)$.

The prior distributions are $f(\theta_1) \sim G(\alpha_{01}, \beta_{01})$, $f(\theta_2) \sim G(\alpha_{02}, \beta_{02})$ and $k \sim \mathcal{U}(0, 1)$, where $G(\alpha, \beta)$ denotes the Gamma distribution with shape parameter α and rate β . We want to approximate the posterior distributions of θ_1 , θ_2 and k using the Gibbs sampling algorithm.

- (a) Plot the data.
- (b) Show that

$$f(\theta_1, \theta_2, k | \mathbf{x}) \propto \theta_1^{\alpha_{01}-1} e^{-\beta_{01}\theta_1} \theta_2^{\alpha_{02}-1} e^{-\beta_{02}\theta_2} \prod_{i=1}^k e^{-\theta_1} \theta_1^{x_i} \prod_{i=k+1}^n e^{-\theta_2} \theta_2^{x_i}$$

and

$$\begin{aligned} \theta_1 | \mathbf{x}, k &\sim G\left(\alpha_{01} + \sum_{i=1}^k x_i, \beta_{01} + k\right) \\ \theta_2 | \mathbf{x}, k &\sim G\left(\alpha_{02} + \sum_{i=k+1}^n x_i, \beta_{02} + n - k\right) \\ f(k | \mathbf{x}, \theta_1, \theta_2) &\propto e^{k(\theta_2 - \theta_1)} \left(\frac{\theta_1}{\theta_2}\right)^{\sum_{i=1}^k x_i}. \end{aligned}$$

- (c) Construct a Gibbs algorithm for this problem. Apply it to the data with $\alpha_{01} = \alpha_{02} = 0.5$ and $\beta_{01} = \beta_{02} = 1$.
- (d) For each parameter, plot the sample path, the histogram of simulated values and the autocorrelation function (use function `acf`).
- (e) Estimate the conditional expectations of each parameter with the associated standard error using the batch means method.