

Computational statistics

Lecture 2: Optimizing smooth multivariate functions

Thierry Denœux

February 23, 2016



Contents of the course (Part I)

- 1 Optimizing smooth univariate functions: Bisection, Newton's method, Fisher scoring, secant method
- 2 Optimizing smooth multivariate functions: nonlinear Gauss-Seidel iteration, Newton's method, Fisher scoring, Gauss-Newton method, ascent algorithms, discrete Newton method, quasi-Newton methods
- 3 Combinatorial optimization: local search, ascent algorithms, simulated annealing, genetic algorithms
- 4 Expectation-Maximization (EM) algorithm for maximizing the likelihood or posterior density



Multivariate optimization for smooth g

- Let $g : \mathbf{x} \in \mathbb{R}^p \rightarrow \mathbb{R}^1$
- Can use analogous stopping criteria:

$$D(\mathbf{x}^{(t+1)}, \mathbf{x}^{(t)}) < \epsilon, \quad \frac{D(\mathbf{x}^{(t+1)}, \mathbf{x}^{(t)})}{D(\mathbf{x}^{(t)}, \mathbf{0})} < \epsilon,$$

or

$$\frac{D(\mathbf{x}^{(t+1)}, \mathbf{x}^{(t)})}{D(\mathbf{x}^{(t)}, \mathbf{0}) + \epsilon} < \epsilon.$$

for $D(\mathbf{u}, \mathbf{v}) = \sum_{i=1}^p |u_i - v_i|$ or $D(\mathbf{u}, \mathbf{v}) = \sqrt{\sum_{i=1}^p (u_i - v_i)^2}$.

- Same strategy of iterative approximation. We will extend previous methods and introduce new options.



Overview

Cyclic coordinate ascent

Newton and quasi-Newton methods

Gauss-Newton method

Nelder-Mead algorithm



Cyclic coordinate ascent

- Also called *backfitting* or *Gauss-Seidel iteration*. One key application is for fitting additive models, GAMs, etc.
- Idea: transform a p -dimensional optimization problem into p univariate optimization problems. How ?




Cyclic coordinate ascent

- Also called *backfitting* or *Gauss-Seidel iteration*. One key application is for fitting additive models, GAMs, etc.
- Idea: transform a p -dimensional optimization problem into p univariate optimization problems. How ?
- Approach: optimize g with respect to each component of \mathbf{x} successively, fixing all other components to their last obtained value.



Algorithm



Case $p = 2$:

- Initialize $x_1 = x_1^{(0)}$
- Find $x_2^{(1)} = \arg \max_{x_2} g(x_1^{(0)}, x_2)$
- Find $x_1^{(1)} = \arg \max_{x_1} g(x_1, x_2^{(1)})$
- Find $x_2^{(2)} = \arg \max_{x_2} g(x_1^{(1)}, x_2)$
- \vdots



Cyclic coordinate ascent: pros and cons

- Advantages:
 - 1 Simplifies a potentially difficult problem
 - 2 Solution of each univariate problem is easier and more stable
- Drawbacks
 - 1 Convergence is not guaranteed
 - 2 Can be slow
- For hard problems (high dimension, complex function shape), we need more sophisticated optimization procedures.



Overview

Cyclic coordinate ascent

Newton and quasi-Newton methods

Gauss-Newton method

Nelder-Mead algorithm



Multivariate Newton's method and Fisher scoring

Multivariate Taylor series argument leads to the following updates:

- Newton's method:

$$\mathbf{x}^{(t+1)} = \mathbf{x}^{(t)} - \mathbf{g}''(\mathbf{x}^{(t)})^{-1} \mathbf{g}'(\mathbf{x}^{(t)}),$$

where


- $\mathbf{g}'(\mathbf{x}) = \left(\frac{\partial g(\mathbf{x})}{\partial x_1}, \dots, \frac{\partial g(\mathbf{x})}{\partial x_p} \right)^T$ is the gradient of g at \mathbf{x}
- $\mathbf{g}''(\mathbf{x}) = \left(\frac{\partial^2 g(\mathbf{x})}{\partial x_i \partial x_j} \right)$ is the $p \times p$ Hessian matrix of g at \mathbf{x}
- Fisher scoring:

$$\boldsymbol{\theta}^{(t+1)} = \boldsymbol{\theta}^{(t)} + \mathbf{I}(\boldsymbol{\theta}^{(t)})^{-1} \boldsymbol{\ell}'(\boldsymbol{\theta}^{(t)}),$$

where $\mathbf{I}(\boldsymbol{\theta}) = -\mathbb{E}\{\boldsymbol{\ell}''(\boldsymbol{\theta})\}$ is the Fisher information matrix at $\boldsymbol{\theta}$.



Example



Two runs starting from $x_a^{(0)}$ and $x_b^{(0)}$ are shown. These converge to the true maximum and to a local minimum, respectively.

Newton's method is not guaranteed to walk uphill. It is not guaranteed to find a local maximum. Step length matters even when step direction is good



Newton-like methods

- Some very effective methods rely on updating equations of the form

$$\mathbf{x}^{(t+1)} = \mathbf{x}^{(t)} - (\mathbf{M}^{(t)})^{-1} \mathbf{g}'(\mathbf{x}^{(t)})$$

where $\mathbf{M}^{(t)}$ is a $p \times p$ matrix approximating the Hessian, $\mathbf{g}''(\mathbf{x}^{(t)})$.

- Two issues:
 - We want to avoid calculating Hessian if it is computationally expensive or analytically difficult
 - We want to guarantee uphill steps



Ascent algorithms

- If we use the updating increment

$$\mathbf{h}^{(t)} = -\alpha^{(t)} [\mathbf{M}^{(t)}]^{-1} \mathbf{g}'(\mathbf{x}^{(t)}).$$

then any positive definite matrix $-\mathbf{M}^{(t)}$ will ensure ascent for a sufficiently small $\alpha^{(t)}$

- **Steepest ascent:** $\mathbf{M}^{(t)} = -\mathbf{I}$. Thus $\mathbf{x}^{(t+1)} = \mathbf{x}^{(t)} + \alpha^{(t)} \mathbf{g}'(\mathbf{x}^{(t)})$
- **Backtracking:** Attempt a step for, say, $\alpha = 1$. If it is downhill, reduce (e.g., halve) α . If the step is still downhill, continue halving $\alpha^{(t)}$ until a sufficiently small step is found to be uphill.
- Notes:
 - $-\mathbf{M}^{(t)} = \mathbf{I}$ is positive definite, therefore backtracking with steepest ascent guarantees uphill steps.
 - $-\mathbf{M}^{(t)} = \mathbf{I}(\theta^{(t)})$ is positive semi-definite. Therefore backtracking with Fisher scoring avoids stepping downhill.



Proof that backtracking works

- For any fixed $\mathbf{x}^{(t)}$ and negative definite $\mathbf{M}^{(t)}$, note that as $\alpha^{(t)} \rightarrow 0$ we have

$$\begin{aligned}g(\mathbf{x}^{(t+1)}) - g(\mathbf{x}^{(t)}) &= g(\mathbf{x}^{(t)} + \mathbf{h}^{(t)}) - g(\mathbf{x}^{(t)}) \\ &= -\alpha^{(t)} \mathbf{g}'(\mathbf{x}^{(t)})^T (\mathbf{M}^{(t)})^{-1} \mathbf{g}'(\mathbf{x}^{(t)}) + o(\alpha^{(t)}),\end{aligned}\tag{1}$$

where the second equality follows from the linear Taylor expansion

$$g(\mathbf{x}^{(t)} + \mathbf{h}^{(t)}) = g(\mathbf{x}^{(t)}) + \mathbf{g}'(\mathbf{x}^{(t)})^T \mathbf{h}^{(t)} + o(\alpha^{(t)}).$$


- Therefore if $-\mathbf{M}^{(t)}$ is positive definite, ascent can be ensured by choosing $\alpha^{(t)}$ sufficiently small, yielding

$$g(\mathbf{x}^{(t+1)}) - g(\mathbf{x}^{(t)}) > 0$$

from (1) since $o(\alpha^{(t)})/\alpha^{(t)} \rightarrow 0$ as $\alpha^{(t)} \rightarrow 0$.



Example



Steepest ascent with backtracking, using $\alpha = 0.25$ initially at each step
The steepest ascent direction is not necessarily the wisest, and
backtracking doesn't prevent oversteps



Discrete Newton method

- To avoid calculating the Hessian, one could resort to an analogue of the 1-dimensional secant method.
- For example,

$$\mathbf{M}_{ij}^{(t)} = \frac{g'_i(\mathbf{x}^{(t)} + h_{ij}^{(t)} \mathbf{e}_j) - g'_i(\mathbf{x}^{(t)})}{h_{ij}^{(t)}}$$

where $g'_i(\mathbf{x}) = dg(\mathbf{x})/dx_i$ is the i th element of $\mathbf{g}'(\mathbf{x})$, \mathbf{e}_j is the p -vector with a 1 in the j th position and zeros elsewhere, and $h_{ij}^{(t)}$ are some constants.

- $h_{ij}^{(t)} = h$ for all (i, j) and t leads to linear convergence order: $\beta = 1$.
- Alternatively, $h_{ij}^{(t)} = x_j^{(t)} - x_j^{(t-1)}$ for all i gives superlinear convergence.



Quasi-Newton methods

- The discrete Newton method strategy is computationally burdensome because $\mathbf{M}^{(t)}$ is wholly updated at every step.
- A more efficient approach can be designed based on the direction of the most recent step. When $\mathbf{x}^{(t)}$ is updated to $\mathbf{x}^{(t+1)} = \mathbf{x}^{(t)} + \mathbf{h}^{(t)}$, the opportunity is presented to learn about the curvature of \mathbf{g}' in the direction of $\mathbf{h}^{(t)}$ near $\mathbf{x}^{(t)}$. information.

- From a first order Taylor series expansion of \mathbf{g}' at $\mathbf{x}^{(t)}$, we have

$$\mathbf{g}'(\mathbf{x}^{(t+1)}) - \mathbf{g}'(\mathbf{x}^{(t)}) \approx \mathbf{g}''(\mathbf{x}^{(t)})(\mathbf{x}^{(t+1)} - \mathbf{x}^{(t)})$$

- $\mathbf{M}^{(t+1)}$ satisfies the secant condition if

$$\mathbf{g}'(\mathbf{x}^{(t+1)}) - \mathbf{g}'(\mathbf{x}^{(t)}) = \mathbf{M}^{(t+1)}(\mathbf{x}^{(t+1)} - \mathbf{x}^{(t)}). \quad (2)$$

- Goal: generate $\mathbf{M}^{(t+1)}$ from $\mathbf{M}^{(t)}$ in a manner that requires few calculations and satisfies (2), while learning about the curvature of \mathbf{g}' in the direction of the most recent step.



Davidon's and BFGS methods

- “Davidon’s update” is the simplest method that meets these requirements. Let $\mathbf{z}^{(t)} = \mathbf{x}^{(t+1)} - \mathbf{x}^{(t)}$ and $\mathbf{y}^{(t)} = \mathbf{g}'(\mathbf{x}^{(t+1)}) - \mathbf{g}'(\mathbf{x}^{(t)})$. Then we can write the update to $\mathbf{M}^{(t)}$ as

$$\mathbf{M}^{(t+1)} = \mathbf{M}^{(t)} + c^{(t)} \mathbf{v}^{(t)} (\mathbf{v}^{(t)})^T$$


where $\mathbf{v}^{(t)} = \mathbf{y}^{(t)} - \mathbf{M}^{(t)} \mathbf{z}^{(t)}$ and $c^{(t)} = \frac{1}{(\mathbf{v}^{(t)})^T \mathbf{z}^{(t)}}$.

- Another well-known method is the “BFGS update”

$$\mathbf{M}^{(t+1)} = \mathbf{M}^{(t)} - \frac{\mathbf{M}^{(t)} \mathbf{z}^{(t)} (\mathbf{M}^{(t)} \mathbf{z}^{(t)})^T}{(\mathbf{z}^{(t)})^T \mathbf{M}^{(t)} \mathbf{z}^{(t)}} + \frac{\mathbf{y}^{(t)} (\mathbf{y}^{(t)})^T}{(\mathbf{z}^{(t)})^T \mathbf{y}^{(t)}}$$

- Davidon’s update does not confer **hereditary positive definiteness**; BFGS does.
- Backtracking is normally used. Monitoring of $c^{(t)}$ is required to maintain numerical stability.

Example



Quasi-Newton optimization with the BFGS update and backtracking to ensure ascent.

Convergence of quasi-Newton methods is generally superlinear, but not quadratic. These are powerful and popular methods, available, for example, in the R function `optim()`.



Overview

Cyclic coordinate ascent

Newton and quasi-Newton methods

Gauss-Newton method

Nelder-Mead algorithm



Gauss-Newton method

Basic idea

- For *nonlinear least squares* problems with observed data (y_i, \mathbf{z}_i) for $i = 1, \dots, n$ and model

$$Y_i = f(\mathbf{z}_i, \boldsymbol{\theta}) + \epsilon_i$$

for some non-linear function, f , and random error, ϵ_i .

- We seek to estimate $\boldsymbol{\theta}$ by maximizing an objective function

$$g(\boldsymbol{\theta}) = - \sum_{i=1}^n (y_i - f(\mathbf{z}_i, \boldsymbol{\theta}))^2.$$

- Newton's method approximates g via Taylor series. The Gauss-Newton approach approximates f itself by its linear Taylor series expansion about $\boldsymbol{\theta}^{(t)}$, so

$$f(\mathbf{z}_i, \boldsymbol{\theta}) \approx f(\mathbf{z}_i, \boldsymbol{\theta}^{(t)}) + (\boldsymbol{\theta} - \boldsymbol{\theta}^{(t)})^T \mathbf{f}'(\mathbf{z}_i, \boldsymbol{\theta}^{(t)})$$

where for each i , $\mathbf{f}'(\mathbf{z}_i, \boldsymbol{\theta}^{(t)})$ is the column vector of partial derivatives of f with respect to $\theta_j^{(t)}$, for $j = 1, \dots, p$, evaluated at $(\mathbf{z}_i, \boldsymbol{\theta}^{(t)})$.



Gauss-Newton method

Linearized reformulation

- Now, instead of $g(\boldsymbol{\theta})$, we maximize

$$\begin{aligned}\tilde{g}(\boldsymbol{\theta}) &= - \sum_{i=1}^n \left(y_i - f(\mathbf{z}_i, \boldsymbol{\theta}^{(t)}) + (\boldsymbol{\theta} - \boldsymbol{\theta}^{(t)})^T \mathbf{f}'(\mathbf{z}_i, \boldsymbol{\theta}^{(t)}) \right)^2 \\ &= - \sum_{i=1}^n \left(x_i^{(t)} - (\boldsymbol{\theta} - \boldsymbol{\theta}^{(t)})^T \mathbf{a}_i^{(t)} \right)^2\end{aligned}$$

with respect to $\boldsymbol{\theta}$, with $x_i^{(t)} = y_i - f(\mathbf{z}_i, \boldsymbol{\theta}^{(t)})$, and define $\mathbf{a}_i^{(t)} = \mathbf{f}'(\mathbf{z}_i, \boldsymbol{\theta}^{(t)})$.

- Then the approximated problem can be re-expressed as minimizing the squared residuals of the linear regression model

$$\mathbf{X}^{(t)} = \mathbf{A}^{(t)}(\boldsymbol{\theta} - \boldsymbol{\theta}^{(t)}) + \boldsymbol{\epsilon}$$

where $\mathbf{X}^{(t)}$ and $\boldsymbol{\epsilon}$ are column vectors whose i th elements consist of $X_i^{(t)}$ and ϵ_i , respectively. Similarly, $\mathbf{A}^{(t)}$ is a matrix whose i th row is $(\mathbf{a}_i^{(t)})^T$.



Gauss-Newton method

Update equation

- This is a linear regression problem! Thus,

$$\boldsymbol{\theta}^{(t+1)} = \boldsymbol{\theta}^{(t)} + \left((\mathbf{A}^{(t)})^T \mathbf{A}^{(t)} \right)^{-1} (\mathbf{A}^{(t)})^T \mathbf{x}^{(t)}.$$

- Requires no computation of Hessian.
- Works best when the model fits fairly well and f is not severely nonlinear.



Overview

Cyclic coordinate ascent

Newton and quasi-Newton methods

Gauss-Newton method

Nelder-Mead algorithm



Nelder-Mead algorithm


Main idea

- An algorithm that does not require the calculation of $g(\mathbf{x})$ or $g''(\mathbf{x})$.
- Idea: evaluation g at $p + 1$ points $\mathbf{x}_1, \dots, \mathbf{x}_p$ forming a **simplex**.
- This simplex defines a region, which is iteratively reshaped by replacing the worst point (vertex) by a better one.



Nelder-Mead algorithm

Definitions




Let

- \mathbf{x}_{best} : vertex with highest value of g
- \mathbf{x}_{worst} : vertex with lowest value of g
- \mathbf{x}_{bad} : 2nd worst vertex
- Best face: face opposite to \mathbf{x}_{worst} , \mathbf{c} its centroid.

Nelder-Mead algorithm

Transformations of a vertex

Five possible transformations of a vertex:



Nelder-Mead algorithm

Basic algorithm

- The location of the new vertex (replacing \mathbf{x}_{worst}) is based on the reflection vertex $\mathbf{x}_r = \mathbf{c} + \alpha_r(\mathbf{c} - \mathbf{x}_{worst})$, usually $\alpha_r = 1$
- If $g(\mathbf{x}_{bad}) < g(\mathbf{x}_r) < g(\mathbf{x}_{best})$: **keep \mathbf{x}_r** as the new vertex
- If $g(\mathbf{x}_r) > g(\mathbf{x}_{best})$: try an **expansion** step
- If $g(\mathbf{x}_r) < g(\mathbf{x}_{bad})$: try a **contraction** step



Nelder-Mead algorithm

Expansion, contraction, shrinking

- **Expansion:** let $\mathbf{x}_e = \mathbf{c} + \alpha_e(\mathbf{x}_r - \mathbf{c})$, usually $\alpha_e = 2$
 - If $g(\mathbf{x}_e) > g(\mathbf{x}_r)$: set \mathbf{x}_e as the new vertex
 - Otherwise, keep \mathbf{x}_r
- **Contraction:**
 - If $g(\mathbf{x}_r) > g(\mathbf{x}_{worst})$: outer contraction. Let $\mathbf{x}_o = \mathbf{c} + \alpha_c(\mathbf{x}_r - \mathbf{c})$, usually $\alpha_c = 0.5$.
 - If $g(\mathbf{x}_o) > g(\mathbf{x}_r)$: keep \mathbf{x}_o
 - Otherwise: perform a shrink transformation
 - If $g(\mathbf{x}_r) \leq g(\mathbf{x}_{worst})$: inner contraction. Let $\mathbf{x}_i = \mathbf{c} + \alpha_c(\mathbf{x}_{worst} - \mathbf{c})$.
 - If $g(\mathbf{x}_i) > g(\mathbf{x}_{worst})$: keep \mathbf{x}_i
 - Otherwise: perform a shrink transformation
- **Shrink transformation:** all vertices except \mathbf{x}_{best} are shrunk toward \mathbf{x}_{best} :



$$\mathbf{x}'_j = \mathbf{x}_{best} + \alpha_s(\mathbf{x}_j - \mathbf{x}_{best}),$$

usually $\alpha_s = 0.5$.



Nelder-Mead algorithm

Example



Nelder-Mead algorithm

Example

